

Why 0.05 BAC Laws Save Lives

Preventing Excessive Drinking is Key

According to the National Transportation Safety Board (NTSB), drivers with a Blood Alcohol Content between 0.05 and 0.079 are seven times more likely to be in a fatal crash than non-drinkers.¹ Nebraska ranks as the second-worst state for self-reported drinking and driving.²

Excessive drinking is a serious and ever-growing problem across the country, but particularly in Nebraska. Excessive consumption is a broad category that includes many types of risky drinking behaviors including:


- Binge drinking – 4+ drinks on one occasion for females, 5+ for males
- Heavy drinking – 8+ drinks per week on average for females, 15+ for males
- Underage drinking – Any drinking by those under the legal drinking age of 21
- Drinking by pregnant women – Consuming any alcohol while pregnant.³

Excessive drinking is a dangerous behavior that is associated with numerous health and social consequences, including but not limited to alcohol-impaired driving, violence, reproductive risks, chronic diseases, and at least seven types of cancer.⁴

The Facts About Driving Impaired

Progress in reducing alcohol-impaired driving has stalled over the past 15 years (see graph below courtesy of the National Academies of Science, Engineering, & Medicine), so it is time to once again consider lowering the legal Blood Alcohol Content (BAC) limits.⁵ Countries around the world have enacted lower BAC laws because the risk of being in a crash increases significantly starting at 0.05 BAC.⁶ Those drivers involved in a fatal crash with a BAC over 0.08 are six times more likely to have a prior conviction for driving while impaired.⁷ Studies have also shown that someone can drive impaired between 200-2,000 times before being arrested.⁸

Alcohol-Impaired Driving Fatalities in the United States, 1982-2016



Number of Fatalities

The plateauing fatality rates indicate that progress has stagnated and even reversed.

9

¹ NTSB (2019)

² Jewett et al. (2015)

³ CDC (2025)

⁴ CDC (2025)

⁵ Fell & Voas (2014)

⁶ Fell (2016)

⁷ NHTSA (2023)

⁸ White & Gasperin (2007)

⁹ NASEM (2018)

It is important to remember that driving while impaired affects not only the impaired driver's safety, but also that of innocent motorists and pedestrians. Indeed, nearly 40% of alcohol-impaired driving fatalities are victims other than the drinking driver.¹⁰

.02 BAC*	About 2 alcoholic drinks**	<ul style="list-style-type: none"> • Decline in visual functions • Decline in ability to perform two tasks at same time
.05 BAC*	About 3 alcoholic drinks**	<ul style="list-style-type: none"> • Reduced coordination • Reduced ability to track moving objects • Difficulty steering • Reduced response to emergency driving situations
.08 BAC*	About 4 alcoholic drinks**	<ul style="list-style-type: none"> • Reduced ability to concentrate • Short-term memory loss • Difficulty controlling speed • Reduced information processing capability • Impaired perception

*Blood Alcohol Concentration measurement. **The number of drinks represents the approximate amount of alcohol that a 160-pound man would need to drink in one hour to reach the listed BAC in each category.

11

The Nebraska Experience

BAC limits in Nebraska:

0.02 BAC – “zero tolerance” laws for individuals under the legal drinking age of 21

0.04 BAC – commercial drivers

0.08 BAC – “per se” laws for all adult drivers on Nebraska roads

Nebraska ranks as the second-worst drinking and driving state with 955 episodes per 1,000 population. Nebraska and Hawaii were the only two states called out by the CDC as having significantly higher rates than the national average.¹²

Between 2014 and 2023, the National Highway Traffic Safety Administration (NHTSA) reports that 796 people were killed in an alcohol-involved crash in Nebraska. Over the past five years, an annual average of 67 alcohol-related traffic fatalities occurred due to an alcohol-impaired driver.¹³

Impacts of 0.05 BAC Laws

When the BAC laws were decreased from 0.1 to 0.08, a 10.4% reduction was seen in alcohol-related fatalities. Reducing the BAC again is estimated to reduce fatal alcohol crashes by another 11% and save at least 1,700 lives annually.¹⁴

According to a 2018 World Health Organization Report, 97 countries around the world had an established BAC limit at or below 0.05%.¹⁵ Some of these countries include Australia, France, Germany, Italy, South Africa, and Turkey. Some countries, including Japan, Norway, Russia, and Sweden, have a BAC limit of 0.02.¹⁶

Implementing 0.05 BAC laws is widely supported by national and international organizations including, but not limited to, the National Transportation Safety Board; National Safety Council; Advocates for Auto and Highway Safety; National Academies of Science, Engineering, and Medicine; World Health Organization; European Transport Safety Council; Canadian Medical Association; 63% of American citizens; and many other national and international bodies.^{17, 18}

¹⁰ NASEM (2018)

¹¹ NTSB (2019)

¹² Jewett et al. (2015)

¹³ NHTSA (2014-2023)

¹⁴ NTSB (2019)

¹⁵ WHO (2018)

¹⁶ Fell & Voas (2014)

¹⁷ NTSB (2019)

¹⁸ Fell & Voas (2006)

Reducing Alcohol-Related Crashes: What Else Works?

In addition to urging states to adopt 0.05% BAC per se laws, a 2018 National Academies of Sciences, Engineering, and Medicine report made the following recommendations for reducing alcohol-related crashes and fatalities:

- Significant **increases in alcohol taxes**;
- **Reduction in availability**, including the number of on- and off-premises alcohol outlets and the days and hours of alcohol sales;
- Adoption and/or strengthening of laws and dedication of enforcement resources to **stop illegal alcohol sales**, including sales to already-intoxicated adults and sales to those under 21;
- **Frequent sobriety checkpoints** with widespread publicity to promote awareness of these efforts;
- Health care systems and health insurers covering and facilitating effective evaluation, prevention, and treatment strategies for binge drinking and alcohol use disorders, including **Screening, Brief Intervention, and Referral to Treatment (SBIRT)**.¹⁹

References

Centers for Disease Control and Prevention (CDC). (2025). Alcohol Use and Your Health. Retrieved March 24, 2026 from <https://www.cdc.gov/alcohol/about-alcohol-use/index.html>

Fell, J. C. (2016). The merits of adopting a 0.05 administrative blood alcohol concentration limit for driving. *American journal of public health*, 106(6), 977.

Fell, J. C., & Voas, R. B. (2006). The effectiveness of reducing illegal blood alcohol concentration (BAC) limits for driving: evidence for lowering the limit to .05 BAC. *Journal of safety research*, 37(3), 233-243.

Fell, J. C., & Voas, R. B. (2014). The effectiveness of a 0.05 blood alcohol concentration (bac) limit for driving in the United States. *Addiction*, 109(6), 869-874.

Jewett, A., Shults, R. A., Banerjee, T., & Bergen, G. (2015). Alcohol-impaired driving among adults—United States, 2012. *Morbidity and mortality weekly report*, 64(30), 814.

National Academies of Sciences, Engineering, and Medicine (NASEM). (2018). *Getting to zero alcohol-impaired driving fatalities: A comprehensive approach to a persistent problem*. National Academies Press.

National Highway Traffic Safety Administration (NHTSA). (2023). 2023 Alcohol-Impaired Driving Traffic Safety Facts. Retrieved March 24, 2026 from <https://crashstats.nhtsa.dot.gov/Api/Public/ViewPublication/813713>

National Highway Traffic Safety Administration (NHTSA). (2014-2023). 2014-2023 Alcohol-Impaired Driving Traffic Safety Facts (multiple documents). Retrieved March 24, 2026 from <https://crashstats.nhtsa.dot.gov/#!/PublicationList/7>

National Transportation Safety Board (NTSB). (2019). 0.05 BAC Safety Briefing Facts. Retrieved October 18, 2023 from <https://portal.ct.gov/-/media/DOT/documents/dvisionzero/NTSB-05-SafetyBriefing-March2019.pdf>

White, W. L., & Gasperin, D. L. (2007). The “hard core drinking driver” identification, treatment and community management. *Alcoholism Treatment Quarterly*, 25(3), 113-132.

World Health Organization (WHO). (2018). Global status report on alcohol and health. Retrieved January 30, 2020 from https://www.who.int/substance_abuse/publications/global_alcohol_report/en/

Contact Information

Project Extra Mile
6001 Dodge Street, CEC 228B
Omaha, NE 68182-0600
Ph: (402) 963-9047
Email: info@projectextramile.org

¹⁹ NASEM (2018)