

# The Facts about Binge Drinking

## *Dangerous Underage Drinking with Dangerous Consequences*

Nebraska has historically had higher levels of underage drinking, binge drinking, and alcohol-impaired driving compared to the rest of the nation.<sup>1</sup>

**Binge drinking** is defined as the consumption of five or more alcoholic beverages in one sitting for a man, or four for a woman. Binge drinking is a dangerous drinking pattern that is associated with a number of health and social consequences.

### Why we care about binge drinking

- **Health** – Binge drinking is associated with the following health problems:
  - Unintentional injuries (e.g. car crashes, falls, burns, drowning)
  - Intentional injuries (e.g. firearm injuries, sexual assaults, domestic violence)
  - Alcohol poisoning
  - Sexually transmitted disease
  - Unintended pregnancy
  - Children born with Fetal Alcohol Spectrum Disorders
  - High blood pressure, stroke, and other cardiovascular diseases
  - Cancers of the breast, mouth, throat, esophagus, liver, and colon
  - Liver disease
  - Neurological damage
  - Sexual dysfunction, and
  - Poor control of diabetes.<sup>2</sup>
- **Safety** – Binge drinkers are 14 times more likely to report alcohol-impaired driving than non-binge drinkers.<sup>3</sup>
- **Economics** – Excessive alcohol consumption costs the citizens of Nebraska over \$1.1 billion dollars in 2010, approximately \$491 million of which fall on government. Seventy-five percent of these costs are attributable to binge drinking.<sup>4</sup>

### The Nebraska experience

According to the results from a recent CDC survey on adult alcohol consumption, Nebraska ranks as the 6<sup>th</sup> worst state in the country for its adult binge drinking rates (19.5%).<sup>5</sup> The city of Lincoln ranks as the worst binge drinking city out of 118 cities indexed across the nation. Three more communities, Omaha, Sioux City, and Grand Island rank among the 30 worst cities in the survey.<sup>6</sup> Furthermore, Nebraska ranks as the fifth-worst state nationally in terms of its excessive alcohol consumption rate (21.7%).<sup>7</sup> Alcohol misuse in the state places a significant strain on the healthcare system, the criminal justice system, and the substance misuse treatment system.

<sup>1</sup> Nebraska DHHS (2015)

<sup>2</sup> CDC (2018)

<sup>3</sup> Naimi et al. (2003)

<sup>4</sup> Sacks et al. (2015)

<sup>5</sup> CDC BRFSS (2021)

<sup>6</sup> CDC (2020)

<sup>7</sup> United Health Foundation (2021)

- Binge drinking rates increase dramatically as age increases and approaches the minimum legal drinking age. The tables below outline the impact that age and sex have had on youth binge drinking rates over the past two years.

**Binge Drinking\*, Rates (%) for Nebraska High School Students, 2018**

Age	Male	Female
9 <sup>th</sup> grade	1.9	2.2
10 <sup>th</sup> grade	4.7	7.2
11 <sup>th</sup> grade	5.5	12.4
12 <sup>th</sup> grade	12.2	18.5

*\* Percentage who reported having five or more drinks for men/four or more drinks for women within a couple of hours, on at least one of the 30 days preceding the survey*


Source: Nebraska Youth Risk Behavior Survey (2018)

**Binge Drinking, Rates (%) for Nebraska Young Adults, 2020**

Age	Male	Female
19-20 years	17.0	19.8
21-22 years	36.0	37.6

Source: Nebraska Young Adult Alcohol Opinion Survey (2020)

- Young adults report having the highest rates of current binge drinking in Nebraska.<sup>8</sup> They also reported having consumed the greatest number of drinks on average during their last binge drinking episode. Eighteen to 24-year-olds reported an average of 8.1 drinks, and 25-34 year-olds reported an average of 8.3. The overall average was 7.2.<sup>9</sup>



- Underage drinking cost the citizens of Nebraska \$324.5 million in 2013. These costs include medical care, work loss, and pain and suffering associated with the ramifications of alcohol use. This translates to a cost of \$1,779 for each youth in the state.<sup>10</sup>

<sup>8</sup> Nebraska DHHS (2015)

<sup>9</sup> Nebraska DHHS (2015)

<sup>10</sup> PIRE (2015)

## Evidence-based interventions for preventing binge drinking

- **State-level policies matter.** A state's policy environment has been found to account for a substantial proportion of state-level variation in binge drinking among US states.<sup>11</sup>
- **Limiting alcohol outlet density.** By limiting the number of retail outlets in an area that sell alcohol, a community can limit access as a means of reducing consumption.<sup>12,13</sup>
- **Consistent enforcement of underage drinking laws and alcohol-impaired driving.** States with more stringent alcohol control policies tend to have lower levels of binge drinking among adults and college students.<sup>14,15,16,17</sup>
- **Increasing taxes on alcohol.** Although alcohol excise taxes are often raised for revenue-generating reasons, several studies suggest that higher excise taxes also have an impact on youth consumption and its consequences. Young adults are more responsive to price increases than adults, and higher taxes increase the consumer price of alcohol. Higher taxes on alcohol are associated with less drinking among 16- to 21-year-olds and high school students.<sup>18</sup>

## References

Babor T.F., Caetano, R., Casswell S., et al. (2003). *Alcohol and Public Policy: No Ordinary Commodity*. New York: Oxford University Press.

Centers for Disease Control and Prevention. (2018). Binge Drinking [fact sheet]. Retrieved on May 10, 2018, from <http://www.cdc.gov/alcohol/fact-sheets/binge-drinking.htm>

Centers for Disease Control and Prevention. (2021). BRFSS Prevalence & Trends Data [online]. Retrieved on October 20, 2022 from <http://www.cdc.gov/brfss/brfssprevalence/index.html>

Centers for Disease Control and Prevention. (2020). SMART: City and County Data - Prevalence Data & Data Analysis Tools, Behavioral Risk Factor Surveillance System, data for 2020. Retrieved on June 10, 2022 from <https://chronicdata.cdc.gov/Behavioral-Risk-Factors/Behavioral-Risk-Factors-Selected-Metropolitan-Area/j32a-sa6u>

Community Preventive Services Taskforce. (n.d.). The Community Guide. *Preventing Excessive Alcohol Consumption. Interventions Directed to the General Population*. Atlanta, GA: Centers for Disease Control and Prevention. Retrieved on March 1, 2017 from <https://www.thecommunityguide.org/topic/excessive-alcohol-consumption>

Naimi, T.S., Brewer, R.D., Mokdad, A., Clark, D., Serdula, M.K., Marks, J.S. (2003). Binge drinking among US adults. *JAMA* 289(1):70–75.

Naimi, T. S., Blanchette, J., Nelson, T. F., Nguyen, T., Oussayef, N., Heeren, T. C., ... & Xuan, Z. (2014). A new scale of the US alcohol policy environment and its relationship to binge drinking. *American journal of preventive medicine*, 46(1), 10-16.

National Research Council and Institute of Medicine. (2004). *Reducing Underage Drinking: A Collective Responsibility*. Washington, DC: National Academies Press.

Nebraska Department of Health and Human Services. (2020). Nebraska Young Adult Alcohol Opinion Survey, 2010 – 2020 State Summary Report. Lincoln, NE: Nebraska Department of Health and Human Services, Division of Behavioral Health.

Nebraska Department of Health and Human Services. (2015). Binge Drinking among Nebraska Adults; presentation on November 10, 2015.

Nebraska Department of Health and Human Services. (2018). Nebraska 2018 Youth Risk Behavior Survey Results. Lincoln, NE: Retrieved on April 30, 2020 from <https://bosr.unl.edu/2018SHARP/20182019YRBSData/2018NEH-Summary-Tables.pdf>

Pacific Institute for Research and Evaluation (PIRE). (2015). Underage Drinking in Nebraska [fact sheet].

Sacks, J. J., Gonzales, K. R., Bouchery, E. E., Tomedi, L. E., & Brewer, R. D. (2015). 2010 National and State Costs of Excessive Alcohol Consumption. *American journal of preventive medicine*, 49(5), e73-e79.

U.S. Department of Health and Human Services. (2007). *The Surgeon General's Call to Action to Prevent and Reduce Underage Drinking*. U.S. Department of Health and Human Services, Office of the Surgeon General. Retrieved on August 4, 2011 from <http://www.surgeongeneral.gov/topics/underagedrinking/>

United Health Foundation. (2021). American's Health Rankings, 2021 Annual Report. Available at <https://assets.americashealthrankings.org/app/uploads/americashealthrankings-2021annualreport.pdf>

Xuan, Z., Blanchette, J.G., Nelson, T.F., Nguyen, T.H., Hadland, S.E., Oussayef, N.L., Heeren, T.C., and Naimi, T.S. (2015). "Youth Drinking in the United States: Relationships with Alcohol Policies and Adult Drinking." *Pediatrics* 136 (1). Retrieved on June 19, 2015, from <http://pediatrics.aappublications.org/content/early/2015/05/26/peds.2015-0537.full.pdf+html>

## Contact Information

Project Extra Mile  
6001 Dodge Street, CEC 228B  
Omaha, NE 68182-0600  
Ph: (402) 963-9047  
Email: [info@projectextramile.org](mailto:info@projectextramile.org)

<sup>11</sup> Naimi et al. (2014)

<sup>12</sup> Babor et al. (2003)

<sup>13</sup> Community Preventive Services Taskforce (n.d.)

<sup>14</sup> National Research Council and Institute of Medicine (2004)

<sup>15</sup> Community Preventive Services Taskforce (n.d.)

<sup>16</sup> U.S. Department of Health and Human Services (2007)

<sup>17</sup> Xuan et al. (2015)

<sup>18</sup> Community Preventive Services Taskforce (n.d.)